



Impact of Water Quality and Phytoplankton on Juvenile Vannamei Shrimp Growth in Low-Salinity Ponds

Eny Heriyati¹, Muhammad Bahrul Ilmi², Chrisna Adhi Suryono^{2*}

¹ Marine Science Study Program, Kutai Timur Agricultural College Jl. Soekarno Hatta No 02, East Kalimantan, 75387, Indonesia

² Department of Marine Science, Faculty of Fisheries and Marine Sciences, Diponegoro University, Semarang, Indonesia

Abstract



Article Info

Received: July 17, 2024

Accepted: August 22, 2024

Published: September 30, 2024

Available online: September 30, 2024

Keywords:

Aquaculture

Growth rate

Phytoplankton

Litopenaeus vannamei

Water quality

*Corresponding Author email:
cas8348@gmail.com

This is an open access article under the CC BY-NC-SA license
(<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

The Vannamei shrimp (*Litopenaeus vannamei*) is a key species in global aquaculture, with its growth influenced by various environmental factors and nutritional quality. This study aimed to analyze water quality parameters and phytoplankton density, as well as assess the growth of juvenile Vannamei shrimp in circular ponds with low salinity (5-6 ppt). Shrimp were stocked at a density of 80 individuals per square meter in 5-ton capacity tanks and fed four times daily over a 30-day period. Water quality parameters, including temperature, dissolved oxygen, pH, and light intensity, were monitored bi-daily, revealing fluctuations that impacted shrimp health. Notably, plankton density assessments showed Cyclotella present on day 14 but absent on day 28, while Microcystis and Rhizosolenia were detected on day 28, indicating shifts in the phytoplankton community. The specific growth rate (SGR) of shrimp was measured at 18.69% per day, with initial and final weights recorded at 0.01 ± 0.00 g and 2.72 ± 0.23 g, respectively. This study highlights the importance of maintaining optimal water quality and understanding phytoplankton dynamics for enhancing shrimp growth. Additionally, it explores the potential benefits of algaline supplementation and the effects of low salinity on the growth of juvenile Vannamei shrimp, contributing valuable insights for sustainable aquaculture practices

Copyright ©2024 Journal of Marine Biotechnology and Immunology.

The juvenile stage in the shrimp's life cycle is crucial, as this is when rapid growth and the development of major anatomy occur (Laramore *et al.*, 2001; Lemos and Weissman, 2021; Seidman and Lawrence, 1985). During this stage, shrimp are more vulnerable to environmental changes and pond conditions, so careful maintenance and management are essential to ensure optimal growth and survival. Feed and nutrition are the main factors influencing the growth of juvenile shrimp (Jaffer *et al.*, 2020; Rodríguez-Olague *et al.*, 2021; Wyban *et al.*, 1995; Yu *et al.*, 2020). High-quality feed is vital to meet the nutritional requirements at this stage.

Phytoplankton, which consists of unicellular algae and cyanobacteria, are the primary producers in aquatic ecosystems (Zohary *et al.*, 2014). They convert solar energy into chemical energy through photosynthesis, making them an important food source for zooplankton and shrimp. Furthermore, phytoplankton plays a significant role in the global biogeochemical cycle by absorbing carbon dioxide (CO₂) from the atmosphere and converting it into organic carbon, helping regulate CO₂ concentrations and potentially

1. Introduction

Litopenaeus vannamei, also known as the Pacific white shrimp, is one of the most popular shrimp species in the global aquaculture industry (Liao and Chien, 2011). The life cycle of the *L. vannamei* consists of several stages, including larval, juvenile, and adult phases (Rojo-Arreola *et al.*, 2020). Nutrition and environmental quality greatly influence growth and development during these phases. Therefore, pond managers must ensure optimal environmental conditions, such as water temperature, salinity, dissolved oxygen, and pH, to support healthy growth. Additionally, providing a balanced diet is crucial to meet the nutritional needs of *L. vannamei* at each growth stage. *L. vannamei* shrimp aquaculture industry continues to grow with the increasing global demand for seafood products. The advancement of technology and sustainable farming practices is a key focus to maintain pond productivity and environmental sustainability (Abdelrahman *et al.*, 2019; Ariadi *et al.*, 2019; Menon *et al.*, 2023; Mugwanya *et al.*, 2022).

2.2.5 Specific Growth Rate

Shrimp were randomly chosen from the population, and their weights were measured on day 0 and day 30 of the rearing period with a precision scale. The specific growth rate (SGR) was subsequently calculated using the following formula (Ario and Nursani, 2024; Ridlo *et al.*, 2024):

$$\text{SGR} (\%/\text{day}) = \frac{(\ln W_t - \ln W_0)}{t} \times 100$$

where W_t is the final weight, W_0 is the initial weight, and t is the time in days.

3. Results

3.1 Water quality parameters

The water quality parameters recorded at 8:00 AM and 4:00 PM exhibited variations (Figure 1). At 8:00 AM, dissolved oxygen levels ranged from 5.66 to 9.26 mg L⁻¹ (Figure 1a), pH levels varied from 7.82 to 8.39 (Figure 1b), temperatures were between 27.7 and 29.8°C (Figure 1c), light intensity ranged from 20,700 to 67,200 Lux (Figure 1d), and salinity was maintained at 5 to 6 ppt (Figure 1e). By 4:00 PM, dissolved oxygen levels ranged from 4.59 to 9.54 mg L⁻¹ (Figure 1a), pH levels varied from 7.6 to 9.3 (Figure 1b), temperatures increased to between 30.1 and 32.5°C (Figure 1c), light intensity ranged from 1,506 to 27,400 Lux (Figure 1d), and salinity remained at 5 to 6 ppt (Figure 1e). These results highlight the fluctuations in water quality parameters throughout the day, offering insights into the dynamic characteristics of the aquatic environment.

3.2 Plankton identification and density

Based on Table 1, the plankton density measurements over 14 and 28 days showed that *Cyclotella* appeared only on day 14 with 750.00 ± 23.6 ($\times 10^3$ cell mL⁻¹) and not showed on day 28. On the other hand, day 28 showed *Microcystis* (11.11×10^3 cell mL⁻¹), *Cyclotella* (529.93×10^3 cell mL⁻¹) and *Rhizosolenia* (11.11×10^3 cell mL⁻¹).

3.3 Specific Growth Rate (SGR)

As shown in Table 2, the initial and final weights were 0.01 ± 0.00 and 2.72 ± 0.23 , respectively. The specific growth rate was measured at 18.69% per day.

4. Discussion

Analyzing the relationship between water quality parameters is a crucial aspect of this study on the growth of juvenile vannamei shrimp. Parameters such as temperature, dissolved oxygen, pH, light intensity, and salinity can have a significant impact on the health and growth of shrimp (Azizah and Samaadan, 2024). Maintaining optimal water quality is essential to create a suitable environment for the growth of vannamei shrimp.

The phytoplankton community in the pond plays a vital role in the growth of vannamei shrimp. Phytoplankton serves as the primary natural food source for shrimp, and variations in the type and quantity of phytoplankton can influence the availability of nutrients (Lyu *et al.*, 2021; Mustika *et al.*, 2023). For instance, *Cyclotella*, a genus of diatoms, is known for its high nutritional content, particularly omega-3 fatty acids, which can enhance shrimp health and growth (Abdel-Moez *et al.*, 2024; Nieri *et al.*, 2023). *Rhizosolenia*, another genus of diatoms, also contributes to the nutritional value of the pond ecosystem, as it is rich in essential fatty acids and serves as a high-quality food source (Damsté *et al.*, 2000; Rowland *et al.*, 2001; Villareal, 1988; Yun *et al.*, 2011; Yung *et al.*, 1988). These

influencing climate change (Jahan, 2023; Naselli-Flores and Padisák, 2023). However, some species of phytoplankton can produce toxins that are harmful to aquatic organisms and humans if they accumulate in contaminated seafood. Environmental conditions such as temperature, light, and nutrients affect the abundance and composition of phytoplankton in ecosystems (Jia *et al.*, 2022; Lee *et al.*, 2019; Pradhan *et al.*, 2022).

Water quality and aquatic parameters, such as temperature, pH, dissolved oxygen, salinity, and nutrient content, are crucial factors in maintaining a healthy aquatic ecosystem (Banerjee *et al.*, 2019; Boyd and Pillai, 1985; Menon *et al.*, 2023). Poor water quality can affect the ecosystem's balance and the survival of aquatic organisms, including shrimp and fish. For example, an increase in water temperature can reduce dissolved oxygen levels, while drastic pH changes can cause stress in aquatic organisms (Boyd, 2015; Boyd, 2019; Boyd and McNevin, 2015; Yusoff *et al.*, 2024). Therefore, monitoring water quality is essential to maintain the health of aquatic ecosystems. The purpose of this study is to measure and analyze water quality parameters, calculate and identify the phytoplankton community, and measure the growth of juvenile Vannamei shrimp in round ponds with low salinity.

2. Material and methods

2.1 Materials

The shrimp utilized in this study had an average weight of around 0.01 ± 0.00 g. They were cultured in circular tanks with a 5-ton capacity, stocked at a density of 80 shrimp per square meter for a period of 30 days. During the experiment, the water salinity was consistently maintained at a low range of 5-6 ppt.

2.2 Methods

2.2.1 Experimental Design

The shrimp used in the study were sourced from the CV Hadid Mukti Karya hatchery. They were introduced into circular ponds with a diameter of 3 meters and a water depth of 1 meter (total depth of 1.2 meters) at a stocking density of 80 shrimp per square meter. Prior to stocking, the shrimp were acclimated by floating their transport bags in the ponds, allowing them to gradually adapt to the pond water temperature. Feeding was conducted four times daily at 07:00, 10:00, 13:00, and 16:00.

2.2.2 Water Quality Sampling Procedure

Water quality parameters, such as temperature, salinity, pH, dissolved oxygen (DO), and light intensity, were monitored twice a day at 8:00 AM and 4:00 PM using standardized instruments.

2.2.3 Plankton Density

Plankton density was assessed using a hemocytometer under a microscope. The magnification varied according to the size of the plankton: larger specimens were counted at low magnification, while smaller or more challenging forms were observed at higher magnifications, sometimes with the use of immersion oil for improved precision. When plankton density was low, the entire bottom of the chamber was analyzed to ensure accuracy. In cases of higher density, only specific sections of the chamber bottom, such as several diagonal fields, were counted to account for irregular distributions while maintaining representativeness in the data. This adaptable approach allowed for precise estimation of plankton density based on the characteristics of the sample (LeGresley and McDermott, 2010; Willén, 1976). Identification of phytoplankton species adhered to the guidelines established

diatoms can provide significant benefits to shrimp health

Heriyati et. al. 2024. Impact of Water Quality and Phytoplankton on.....
and growth when present in sufficient quantities.




Figure 1. The water quality parameters. a). dissolved oxygen levels; b) pH levels; c) temperatures were ; d) light intensity; e) salinity.

Tabel 1. Plankton Identification and Density on 14 days and 28 days of rearing

| Days | the plankton density ($\times 10^3$ cell mL ⁻¹) | | | |
|------|--|--------------|----------------|--------------|
| | Cyclotella | Microcystis | Cyclotella | Rhizosolenia |
| 14 | 750.00 ± 23.6 | ND | ND | ND |
| 28 | ND | 11.11 ± 0.00 | 529.93 ± 66.97 | 11.11 ± 0.00 |

ND : Not detected

Table 2. Weight of *Litopaneus vannamei*

| Initial weight (gr) | Final weight (gr) | SGR(%) |
|---------------------|-------------------|--------|
| 0,01±00 | 2.72 ± 0.23 | 18.69% |

On the other hand, *Microcystis*, a genus of cyanobacteria, can have a different impact on shrimp growth. While some cyanobacteria can be beneficial in the food web, *Microcystis* is often associated with harmful algal blooms (HABs) that produce toxins such as microcystins, which can be detrimental to shrimp health (Lad *et al.*, 2022; Le Manach *et al.*, 2019; Le *et al.*, 2023; Wilhelm *et al.*, 2020). The presence of *Microcystis* in the pond environment can affect water quality and potentially inhibit the growth of *vannamei* shrimp, especially if the population density becomes excessive.

This study will investigate how variations in the phytoplankton community, particularly the presence of *Microcystis*, *Cyclotella*, and *Rhizosolenia*, affect the growth of *vannamei* shrimp and how these species relate to changes in water quality parameters.

The use of alginate as a supplement in *vannamei* shrimp farming is another focus of this study. Alginate is a compound known to enhance feed efficiency and improve shrimp health (Azhar and Yudiat, 2023; Chen *et al.*, 2022; Yudiat *et al.*, 2016; Yudiat *et al.*, 2019). This research will evaluate the extent to which alginate supplementation can promote the growth of juvenile *vannamei* shrimp in circular ponds. The findings will provide valuable insights into the potential application of alginate in improving *vannamei* shrimp aquaculture.

This study will also analyze how low salinity affects the growth and health of *vannamei* shrimp. Understanding its impact is essential for effective pond management and ensuring the optimal growth of juvenile *vannamei* shrimp.

This study demonstrates that water quality and phytoplankton community dynamics significantly impact the growth of *Vannamei* shrimp (*Litopenaeus vannamei*). Fluctuations in water quality parameters, such as temperature, dissolved oxygen, pH, and light intensity, influence the health and growth of shrimp during the juvenile phase. The measurements indicate that the presence of *Cyclotella*, *Microcystis*, and *Rhizosolenia* within the pond ecosystem contributes to the variability in available nutrition for the shrimp. The recorded specific growth rate (SGR) of 18.69% per day highlights the potential for substantial growth under well-managed conditions.

Additionally, the study underscores the importance of regular water quality monitoring and phytoplankton management to ensure optimal growth environments for shrimp. The implementation of alginate supplementation could serve as a promising strategy to enhance shrimp health and feed efficiency. These findings provide valuable insights for sustainable aquaculture practices, particularly in managing low salinity levels in ponds. Effective management of these environmental factors is crucial for achieving maximum productivity and sustainability within the aquaculture industry.

3 Conclusions

This study demonstrates that water quality and phytoplankton dynamics significantly influence the growth of *Vannamei* shrimp (*Litopenaeus vannamei*). Key factors such as temperature, dissolved oxygen, pH, and light intensity affect shrimp health, while the presence of *Cyclotella*, *Microcystis*, and *Rhizosolenia* impacts nutritional availability. With a specific growth rate (SGR) of 18.69% per day, optimal conditions are crucial for strong growth.

Ethics approval

No permits were required.

Data availability statement

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Author contributions

MBI is doing research ideas, sample image collection, water quality sampling collection, EH and CAS are supervising and writing

Funding

No funding available

Acknowledgments

I would like to state my special thanks of gratefulness to Diponegoro University.

Declaration of competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

References

Abdel-Moez, A. M., Ali, M. M., El-gandy, G., Mohammady, E. Y., Jarmołowicz, S., El-Haroun, E., Elsaied, H. E., & Hassaan, M. S. 2024. Effect of including dried microalgae *Cyclotella meneghiniana* on the reproductive performance, lipid metabolism profile and immune response of Nile tilapia broodstock and offspring. *Aquaculture Reports*. 36: 102099. <https://doi.org/10.1016/j.agrep.2024.102099>

Abdelrahman, H. A., Abebe, A., & Boyd, C. E. 2019. Influence of variation in water temperature on survival, growth and yield of Pacific white shrimp *Litopenaeus vannamei* in inland ponds for low-salinity culture. *Aquaculture Research*. 50(2): 658-672. <https://doi.org/10.1111/are.13943>

Ariadi, H., Fadjar, M., Mahmudi, M., & Supriatna, S. 2019. The relationships between water quality parameters and the growth rate of white shrimp (*Litopenaeus vannamei*) in intensive ponds. *AACL Bioflux*. 12: 2103-2216.

Ario, R., & Nursani, I. A. 2024. Influence of Water Quality and Phytoplankton Community on the Growth of *Litopenaeus vannamei* in Low-Salinity Semi-Mass Circular Ponds. *Journal of Marine Biotechnology and Immunology*. 2(2): 7-12. 10.61741/7kkn2e98

Azhar, N., & Yudiat, E. 2023. Outbreak simulation of *Litopenaeus vannamei* recovery rate with oral alginate and spirulina diet supplementation against *Vibrio parahaemolyticus* AHPND. *Aquacult Int*. 31(3): 1659-1676. <https://doi.org/10.1007/s10499-023-01050-6>

Azizah, K. N., & Samaadan, G. M. 2024. Analysis of the Relationship between Water Quality Parameters and Phytoplankton Communities on the Growth of *Litopenaeus vannamei* Post-Larvae in Low Salinity

Circular Ponds. *Journal of Marine Biotechnology and Immunology*. 2(1): 24-29. 10.61741/b3fwzr29

Banerjee, A., Chakrabarty, M., Rakshit, N., Bhowmick, A. R., & Ray, S. 2019. Environmental factors as indicators of dissolved oxygen concentration and zooplankton abundance: Deep learning versus traditional regression approach. *Ecological Indicators*. 100: 99-117. <https://doi.org/10.1016/j.ecolind.2018.09.051>

Boyd, C. 2015. *Water Quality: An Introduction*. <https://doi.org/10.1007/978-3-319-17446-4>

Boyd, C., & Pillai, V. 1985. Water Quality Management in Aquaculture.

Boyd, C. E. 2019. *Water quality: an introduction*. Springer Nature.

Boyd, C. E., & McNevin, A. A. (2015). Water Use by Aquaculture Systems. In *Aquaculture, Resource Use, and the Environment* (pp. 101-122). Retrieved 2024/05/27, from <https://doi.org/10.1002/9781118857915.ch6>

Chen, G., Liu, B., Chen, J., Liu, H., Tan, B., Dong, X., Yang, Q., Chi, S., Zhang, S., & Yao, M. 2022. Supplementing Sulfate-Based Alginate Polysaccharide Improves Pacific White Shrimp (*Litopenaeus vannamei*) Fed Fishmeal Replacement with Cottonseed Protein Concentrate: Effects on Growth, Intestinal Health, and Disease Resistance. *Aquaculture Nutrition*. 2022(1): 7132362. <https://doi.org/10.1155/2022/7132362>

Damsté, J. S. S., Schouten, S., Rijpstra, W. I. C., Hopmans, E. C., Peletier, H., Gieskes, W. W. C., & Geenevasen, J. A. J. 2000. Novel polyunsaturated n-alkenes in the marine diatom *Rhizosolenia setigera*. *European Journal of Biochemistry*. 267(18): 5727-5732. <https://doi.org/10.1046/j.1432-1327.2000.01636.x>

Genkal, S. I. 2012. Morphology, taxonomy, ecology, and distribution of *Cyclotella choctawhatcheeana* prasad (Bacillariophyta). *Inland Water Biology*. 5(2): 169-177. <https://doi.org/10.1134/S1995082912020046>

Jaffer, Y. D., Saraswathy, R., Ishfaq, M., Antony, J., Bundela, D. S., & Sharma, P. C. 2020. Effect of low salinity on the growth and survival of juvenile pacific white shrimp, *Penaeus vannamei*: A revival. *Aquaculture*. 515: 734561. <https://doi.org/10.1016/j.aquaculture.2019.734561>

Jahan, S. 2023. The Role of Phytoplanktons in the Environment and in Human Life, a Review. *BASRA JOURNAL OF SCIENCE*. 41: 392-411. 10.29072/basjs.20230212

Jia, J., Chen, Q., Ren, H., Lu, R., He, H., & Gu, P. 2022. Phytoplankton Composition and Their Related Factors in Five Different Lakes in China: Implications for Lake Management. *Int J Environ Res Public Health*. 19(5). 10.3390/ijerph19053135

Kilham, P., & Hecky, R. E. 1988. Comparative ecology of marine and freshwater phytoplankton1. *Limnology and Oceanography*. 33(4part2): 776-795. <https://doi.org/10.4319/lo.1988.33.4part2.0776>

Lad, A., Breidenbach, J. D., Su, R. C., Murray, J., Kuang, R., Mascarenhas, A., Najjar, J., Patel, S., Hegde, P., Youssef, M., Breuler, J., Kleinhenz, A. L., Ault, A. P., Westrick, J. A., Modyanov, N. N., Kennedy, D. J., & Haller, S. T. 2022. As We Drink and Breathe: Adverse Health Effects of Microcystins and Other

Heriyati et. al. 2024. *Impact of Water Quality and Phytoplankton on.....* Harmful Algal Bloom Toxins in the Liver, Gut, Lungs and Beyond. *Life (Basel)*. 12(3). 10.3390/life12030418

Laramore, S., Laramore, C. R., & Scarpa, J. 2001. Effect of Low Salinity on Growth and Survival of Postlarvae and Juvenile *Litopenaeus vannamei*. *Journal of the World Aquaculture Society*. 32(4): 385-392. <https://doi.org/10.1111/j.1749-7345.2001.tb00464.x>

Le Manach, S., Duval, C., Marie, A., Djediat, C., Catherine, A., Edery, M., Bernard, C., & Marie, B. 2019. Global Metabolomic Characterizations of *Microcystis* spp. Highlights Clonal Diversity in Natural Bloom-Forming Populations and Expands Metabolite Structural Diversity. *Front. Microbiol*. 10. 10.3389/fmicb.2019.00791

Le, V. V., Ko, S.-R., Kang, M., Oh, H.-M., & Ahn, C.-Y. 2023. Effective control of harmful *Microcystis* blooms by paucibactin A, a novel macrocyclic tambjamine, isolated from *Paucibacter aquatile* DH15. *Journal of Cleaner Production*. 383: 135408. <https://doi.org/10.1016/j.jclepro.2022.135408>

Lee, K. H., Jeong, H. J., Lee, K., Franks, P., Seong, K., Lee, S. Y., Lee, M. J., Jang, S. H., Potvin, E., Lim, A. S., Yoon, E., Yoo, Y., Kang, N., & Kim, K. Y. 2019. Effects of warming and eutrophication on coastal phytoplankton production. *Harmful Algae*. 81: 106-118. 10.1016/j.hal.2018.11.017

LeGresley, M., & McDermott, G. 2010. Counting chamber methods for quantitative phytoplankton analysis- haemocytometer, Palmer-Malone cell and Sedgewick-Rafter cell. *Microscopic and Molecular Methods for Quantitative Phytoplankton Analysis*: 25-30.

Lemos, D., & Weissman, D. 2021. Moulting in the grow-out of farmed shrimp: a review. *Reviews in Aquaculture*. 13(1): 5-17. <https://doi.org/10.1111/raq.12461>

Liao, I. C., & Chien, Y.-H. (2011). The Pacific White Shrimp, *Litopenaeus vannamei*, in Asia: The World's Most Widely Cultured Alien Crustacean. In B. S. Galil, P. F. Clark, & J. T. Carlton (Eds.), *In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts* (pp. 489-519). Springer Netherlands. https://doi.org/10.1007/978-94-007-0591-3_17 https://doi.org/10.1007/978-94-007-0591-3_17

Lyu, T., Yang, W., Cai, H., Wang, J., Zheng, Z., & Zhu, J. 2021. Phytoplankton community dynamics as a metrics of shrimp healthy farming under intensive cultivation. *Aquaculture Reports*. 21: 100965. <https://doi.org/10.1016/j.aqrep.2021.100965>

Menon, S. V., Kumar, A., Middha, S. K., Paital, B., Mathur, S., Johnson, R., Kademan, A., Usha, T., Hemavathi, K. N., Dayal, S., Ramalingam, N., Subaramaniyam, U., Sahoo, D. K., & Asthana, M. 2023. Water physicochemical factors and oxidative stress physiology in fish, a review. 11. 10.3389/fenvs.2023.1240813

Mugwanya, M., Dawood, M. A. O., Kimera, F., & Sewilam, H. 2022. Anthropogenic temperature fluctuations and their effect on aquaculture: A comprehensive review. *Aquaculture and Fisheries*. 7(3): 223-243. <https://doi.org/10.1016/j.aaf.2021.12.005>

Mustika, P., Ren, F., Kasprijo, & Yasmin, M. 2023. Phytoplankton Community in Vannamei Shrimp (*Litopenaeus vannamei*) Cultivation in Intensive Ponds. *IRAQI JOURNAL OF AGRICULTURAL SCIENCES*. 54(1): 134-146. 10.36103/ijas.v54i1.1684

Naselli-Flores, L., & Padisák, J. 2023. Ecosystem services provided by marine and freshwater phytoplankton. *Hydrobiologia*. 850(12-13): 2691-2706. 10.1007/s10750-022-04795-y

Nieri, P., Carpi, S., Esposito, R., Costantini, M., & Zupo, V. 2023. Bioactive Molecules from Marine Diatoms and Their Value for the Nutraceutical Industry. *Nutrients*. 15(2). 10.3390/nu15020464

Pradhan, B., Kim, H., Abassi, S., & Ki, J. S. 2022. Toxic Effects and Tumor Promotion Activity of Marine Phytoplankton Toxins: A Review. *Toxins (Basel)*. 14(6). 10.3390/toxins14060397

Reynolds, C. S. 1984. The ecology of freshwater phytoplankton.

Ridlo, A., Firdaus, M. L. M., & Sumarwan, J. 2024. Growth of *Litopenaeus vannamei* using Synbiotics Supplementation Diet in Outdoor Low-Salinity Ponds Concerning Water Quality Parameters and Phytoplankton Communities. *Journal of Marine Biotechnology and Immunology*. 2(2): 32-36. 10.61741/d2wnd591

Rodríguez-Olague, D., Ponce-Palafox, J. T., Castillo-Vargasmachuca, S. G., Arámbul-Muñoz, E., de los Santos, R. C., & Esparza-Leal, H. M. 2021. Effect of nursery system and stocking density to produce juveniles of whiteleg shrimp *Litopenaeus vannamei*. *Aquaculture Reports*. 20: 100709. <https://doi.org/10.1016/j.aqrep.2021.100709>

Rojo-Arreola, L., García-Carreño, F., Romero, R., & Díaz Dominguez, L. 2020. Proteolytic profile of larval developmental stages of *Penaeus vannamei*: An activity and mRNA expression approach. *PLoS One*. 15(9): e0239413. 10.1371/journal.pone.0239413

Rowland, S. J., Allard, W. G., Belt, S. T., Massé, G., Robert, J. M., Blackburn, S., Frampton, D., Revill, A. T., & Volkman, J. K. 2001. Factors influencing the distributions of polyunsaturated terpenoids in the diatom, *Rhizosolenia setigera*. *Phytochemistry*. 58(5): 717-728. [https://doi.org/10.1016/S0031-9422\(01\)00318-1](https://doi.org/10.1016/S0031-9422(01)00318-1)

Seidman, E. R., & Lawrence, A. L. 1985. Growth, Feed Digestibility, And Proximate Body Composition Of Juvenile *Penaeus vannamei* And *Penaeus monodon* Grown At Different Dissolved Oxygen Levels. *J World Aquac Soc*. 16(1-4): 333-346. <https://doi.org/10.1111/j.1749-7345.1985.tb00214.x>

Villareal, T. A. 1988. Positive buoyancy in the oceanic diatom *Rhizosolenia debyana* H. Peragallo. *Deep Sea Research Part A. Oceanographic Research* Heriyati et. al. 2024. *Impact of Water Quality and Phytoplankton on... Papers*. 35(6): 1037-1045. [https://doi.org/10.1016/0198-0149\(88\)90075-1](https://doi.org/10.1016/0198-0149(88)90075-1)

Wilhelm, S. W., Bullerjahn, G. S., & McKay, R. M. L. 2020. The Complicated and Confusing Ecology of *Microcystis* Blooms. *mBio*. 11(3). 10.1128/mBio.00529-20

Willén, E. 1976. A simplified method of phytoplankton counting. *British Phycological Journal*. 11(3): 265-278. <https://doi.org/10.1080/00071617600650551>

Wyban, J., Walsh, W. A., & Godin, D. M. 1995. Temperature effects on growth, feeding rate and feed conversion of the Pacific white shrimp (*Penaeus vannamei*). *Aquaculture*. 138(1): 267-279. [https://doi.org/10.1016/0044-8486\(95\)00032-1](https://doi.org/10.1016/0044-8486(95)00032-1)

Yu, Q., Xie, J., Huang, M., Chen, C., Qian, D., Qin, J. G., Chen, L., Jia, Y., & Li, E. 2020. Growth and health responses to a long-term pH stress in Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture Reports*. 16: 100280. <https://doi.org/10.1016/j.aqrep.2020.100280>

Yudiati, E., Isnansetyo, A., Murwantoko, Ayuningtyas, Triyanto, & Handayani, C. R. 2016. Innate immune-stimulating and immune genes up-regulating activities of three types of alginate from *Sargassum siliquosum* in Pacific white shrimp, *Litopenaeus vannamei*. *Fish Shellfish Immunol*. 54: 46. <https://doi.org/10.1016/j.fsi.2016.03.022>

Yudiati, E., Isnansetyo, A., Murwantoko, Triyanto, & Handayani, C. R. 2019. Alginate from *Sargassum siliquosum* simultaneously stimulates innate immunity, upregulates immune genes, and enhances resistance of Pacific white shrimp (*Litopenaeus vannamei*) against white spot syndrome virus (WSSV). *Mar Biotechnol*. 21(4): 503. <https://doi.org/10.1007/s10126-019-09898-7>

Yun, S. M., Lee, S. D., & Lee, J. 2011. Morphology and distribution of some marine diatoms, Family Rhizosoleniaceae, in Korean coastal waters: a genus *Rhizosolenia* 1. *Algae*. 26: 141-152. 10.4490/algae.2011.26.2.141

Yung, Y. K., Nicholls, K. H., & Cheng, A. G. 1988. The detection of *Rhizosolenia* (Bacillariophyceae) in sediment of Ontario lakes and implications for paleoecology. *Journal of Paleolimnology*. 1(1): 61-69. 10.1007/BF00202194

Yusoff, F. M., Umi, W. A. D., Ramli, N. M., & Harun, R. 2024. Water quality management in aquaculture. *Cambridge Prisms: Water*. 2: e8. 10.1017/wat.2024.6

Zohary, T., Yacobi, Y. Z., Alster, A., Fishbein, T., Lippman, S., & Tibor, G. (2014). Phytoplankton. In T. Zohary, A. Sukenik, T. Berman, & A. Nishri (Eds.), *Lake Kinneret: Ecology and Management* (pp. 161-190). Springer Netherlands. https://doi.org/10.1007/978-94-017-8944-8_10 https://doi.org/10.1007/978-94-017-8944-8_10